


# Postoperative Admission of Adult Craniotomy Patients to the Neuroscience Ward Reduces Length of Stay and Cost

Matthew Z. Sun, MD <sup>\*</sup>  
 Diana Babayan, MPH<sup>\*</sup>  
 Jia-Shu Chen, BS<sup>\*</sup>  
 Maxwell M. Wang, BS<sup>\*</sup>  
 Priyanka K. Naik, BS<sup>\*</sup>  
 Kara Reitz, BS<sup>\*</sup>  
 Jingyi Jessica Li, PhD<sup>†</sup>  
 Nader Pouratian, MD, PhD <sup>\*</sup>  
 Won Kim, MD<sup>\*</sup>

<sup>\*</sup>Department of Neurosurgery, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, California, USA; <sup>†</sup>Department of Statistics, University of California Los Angeles, Los Angeles, California, USA

An abstract of this manuscript was presented at the 2019 CNS Annual Meeting in San Francisco, California on October 22, 2019.

## Correspondence:

Won Kim, MD,  
 Department of Neurosurgery,  
 Ronald Reagan UCLA Medical Center,  
 David Geffen School of Medicine at  
 UCLA,  
 300 Stein Plaza Suite 420,  
 Los Angeles, CA 90095, USA.  
 Email: [WonKim@mednet.ucla.edu](mailto:WonKim@mednet.ucla.edu)

Received, December 31, 2019.  
 Accepted, December 13, 2020.

© Congress of Neurological Surgeons 2021. All rights reserved. For permissions, please e-mail: [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

**BACKGROUND:** The neurointensive care unit (NICU) has traditionally been the default recovery unit after elective craniotomies.

**OBJECTIVE:** To assess whether admitting adult patients without significant comorbidities to the neuroscience ward (NW) instead of NICU for recovery resulted in similar clinical outcome while reducing length of stay (LOS) and hospitalization cost.

**METHODS:** We retrospectively analyzed the clinical and cost data of adult patients undergoing supratentorial craniotomy at a university hospital within a 5-yr period who had a LOS less than 7 d. We compared those admitted to the NICU for 1 night of recovery versus those directly admitted to the NW.

**RESULTS:** The NICU and NW groups included 340 and 209 patients, respectively, and were comparable in terms of age, ethnicity, overall health, and expected LOS. NW admissions had shorter LOS (3.046 vs 3.586 d,  $P < .001$ ), and independently predicted shorter LOS in multivariate analysis. While the NICU group had longer surgeries (6.8 vs 6.4 h), there was no statistically significant difference in the cost of surgery. The NW group was associated with reduced hospitalization cost by \$3193 per admission on average ( $P < .001$ ). Clinically, there were no statistically significant differences in the rate of return to Operating Room, Emergency Department readmission, or hospital readmission within 30 d.

**CONCLUSION:** Admitting adult craniotomy patients without significant comorbidities, who are expected to have short LOS, to NW was associated with reduced LOS and total cost of admission, without significant differences in postoperative clinical outcome.

**KEY WORDS:** Craniotomy postop, Neuroscience ward, Length of stay, Admission cost, Clinical outcome

*Neurosurgery* 0:1–9, 2021

DOI:10.1093/neuros/nyab089

[www.neurosurgery-online.com](http://www.neurosurgery-online.com)

A major component of high-cost healthcare originates from patients who require hospitalization after a major surgery, and, in particular, from cost associated with utilization of the intensive care unit (ICU) in the postoperative period.<sup>1,2</sup> Recent data have linked ICU length of stay (LOS) to non-ICU<sup>1</sup> and overall hospital LOS.<sup>2</sup> The neurointensive care unit (NICU) is one of

the costliest units<sup>3</sup> because of higher staff-to-patient ratio and additional neuromonitoring equipment necessary for patient care. Traditionally, patients undergoing craniotomies for tumor resection are admitted to the NICU after the surgery. However, there is no class I evidence demonstrating that patients perform better when they recover in the NICU as opposed to the neuroscience ward (NW). Conversely, the lack of evidence to the contrary limits a neurosurgeon's ability to justify admitting such patients to a non-ICU unit postoperatively.

Our department initiated a clinical pathway years ago that gave our neurosurgeons the flexibility to admit patients without significant medical comorbidities to the NW postoperatively after uncomplicated elective supratentorial craniotomies for brain tumor resection. Here, we present a retrospective cost-effectiveness analysis of our institutional experience over a 5-yr period

**ABBREVIATIONS:** AIC, Akaike Information Criterion; APR-DRG, All Patient Refined Diagnosis Related Group; ED, Emergency Department; LOS, length of stay; MS-DRG, Medicare Severity Diagnosis Related Group; NICU, neurointensive care unit; NW, neuroscience ward; OR, Operating Room; PACU, Post-Anesthesia Care Unit

Supplemental digital content is available for this article at [www.neurosurgery-online.com](http://www.neurosurgery-online.com).

to compare patients who recovered in the NICU vs the NW after elective craniotomy for tumor resection. This is the first time an analysis of a large series of patients, such as this one, has been reported.

## METHODS

### Patient Selection

We included all elective supratentorial craniotomy patients who were 18 yr or older with a LOS of 7 d or less who were admitted to our university hospital from March 2013 to April 2018. We excluded patients who underwent surgical procedures for shunts, Ommaya reservoirs, hemispherectomies, endoscopic transnasal approaches, external ventricular drains, stereotactic burr hole craniotomies, vascular and interventional radiology procedures, and any posterior fossa pathology. The decision to admit to the NW vs NICU was made by surgeon's practice preference.

### Data Collection

An institutional review board approval exempted us from patient consent because only de-identified information was collected for the purposes of quality improvement. We identified admissions based on a combination of International Classification of Diseases codes and procedure performed. The complete list of ICD9 codes is presented in **Supplemental Digital Content 1, Table 1**. The complete list of variables is listed in **Supplemental Digital Content 2, Table 2**. The variables analyzed are notable for patient demographics, financial information, admission data, predicting variables, LOS data, cost data, and any confounding variables. Medicare Severity Diagnosis Related Groups (MS-DRGs), MS-DRG weight, and All Patient Refined Diagnosis Related Group (APR-DRG) were collected for each admission to control for severity of illness and any comorbidities. The hospital financial decision team provided the calculated expected LOS based on the Vizient Clinical Data Base/Resource Manager model (Vizient Inc, Texas).


### Statistical Analysis

The RStudio (RStudio Inc, Version 1.1.453) software was used to perform statistical analyses. Qualitative and quantitative outcome variables between the 2 groups were compared with the Pearson's chi-squared test and the Wilcoxon-Mann-Whitney test, respectively. Multivariate linear and logistic regression models were constructed, starting with univariate regressions between the outcomes and every individual covariate to identify potential predictors for each outcome. Any variable with a  $P$ -value less than .10 was included in a bidirectional stepwise regression for each outcome. Through each stepwise model selection, the regression model with the lowest Akaike Information Criterion (AIC) was selected and used to identify and evaluate independent predictors of the outcomes of interest. The  $P$ -value for statistical significance was set at .05.

## RESULTS

### Admitting to the NW Is Associated With Reduced Overall LOS

Our analysis was limited to adult patients without significant comorbidities who were not expected to need prolonged recovery



in the NICU. We retrospectively analyzed all patients 18 yr or older who underwent elective supratentorial craniotomy at our university's main medical center from March 2013 to March 2018, were hospitalized for 7 d or less, and stayed in the NICU for either 0 or 1 d, and assigned them to the NW and NICU groups, respectively. While most patients recover for at least 2 h in the Post-Anesthesia Care Unit (PACU) before transferring to either the NW or the NICU, some were directly admitted to the NICU from the Operating Room (OR) (Figure 1).

A total of 209 patients were admitted to the NW, while 340 were admitted to the NICU. The patient characteristics that may be potential confounders were mostly comparable between the NW and NICU groups (Table 1), including age, MS-DRG, APR-DRG severity, ethnicity, insurance type, and cost of surgery. However, the NICU group had longer OR time usage, higher percentage of male patients, different discharge disposition, MS-DRG weight, and day of admission grouped by days of the week, when compared to the NW group.

The average LOS for the NW group was 3.046 d, compared to 3.586 d for the NICU group ( $P < .001$ ) (Table 2). There was no difference in the expected LOS between the NW and NICU groups ( $P = .184$ ) (Table 2, Table 3A).

In our multivariate analysis, postoperative stay in the NICU even for 1 night was independently associated with increased LOS ( $P = .015$ ), but not with expected LOS (Table 3B). Additionally, MS-DRG weight, OR hours charged, day of admission, discharge disposition, and APR-DRG severity also independently predicted increased LOS (Table 3B).

Further subgroup analysis of those who stayed in the PACU past midnight on postoperative day 0 showed that it was

**TABLE 1. Patient Demographics and Univariate Analysis of Controlled Variables**

	NW (n = 209)	NICU (n = 340)	
<b>Continuous variables</b>	<b>Average; 95% CI</b>		<b>Mann-Whitney P value</b>
Age at discharge	52.67; [50.76, 54.57]	51.19; [49.63, 52.74]	.208
MS DRG weight	2.97; [2.85, 3.09]	3.15; [3.05, 3.26]	.048*
OR hours charged	6.43; [6.14, 6.72]	6.83; [6.64, 7.01]	<.001***
<b>Categorical variables</b>	<b>Frequency</b>		<b>Pearson's chi-squared P value</b>
<b>Sex</b>			.038*
Female	115 (55.0%)	156 (45.9%)	
Male	94 (45.0%)	184 (54.1%)	
<b>Ethnicity</b>			.471
Native American	0	1 (0.3%)	
African American	6 (2.9%)	13 (3.8%)	
Asian	26 (12.4%)	34 (10.0%)	
Caucasian	142 (67.9%)	249 (73.2%)	
Pacific Islander	0	1 (0.3%)	
Other	35 (16.7%)	42 (12.4%)	
<b>Insurance</b>			.461
Commercial	136 (65.1%)	209 (61.5%)	
Group Health Plan	1 (0.5%)	4 (1.2%)	
International	0	3 (0.9%)	
Medi-Cal	5 (2.4%)	6 (1.8%)	
Medicare	42 (20.1%)	79 (23.2%)	
Self-pay	1 (0.5%)	6 (1.8%)	
Tricare	2 (1.0%)	6 (1.8%)	
Managed care	11 (5.3%)	17 (5.0%)	
Worker's comp	1 (0.5%)	0	
Other	10 (4.8%)	10 (2.9%)	
<b>Day of admission</b>			.019*
Monday	44 (21.1%)	72 (21.2%)	
Tuesday	46 (22.0%)	106 (31.2%)	
Wednesday	22 (10.5%)	16 (4.7%)	
Thursday	61 (29.2%)	103 (30.3%)	
Friday	35 (16.7%)	43 (12.6%)	
Saturday	1 (0.5%)	0	
<b>Discharge disposition</b>			.005*
Acute care	0	4 (1.2%)	
Deceased	0	1 (0.3%)	
Home	186 (89.0%)	262 (77.1%)	
Inpatient rehab	22 (10.5%)	62 (18.2%)	
Psychiatric hospital	0	0	
Skilled nursing facility	1 (0.5%)	11 (3.2%)	
<b>MS DRG</b>			.133
23.00	4 (1.9%)	7 (2.1%)	
25.00	48 (23.0%)	110 (32.4%)	
26.00	49 (23.4%)	73 (21.5%)	
27.00	108 (51.7%)	143 (42.1%)	
41.00	0	3 (0.9%)	
42.00	0	1 (0.3%)	
54.00	0	2 (0.6%)	
55.00	0	1 (0.3%)	
<b>APR DRG severity</b>			.125
Minor	77 (36.8%)	100 (29.4%)	
Moderate	109 (52.2%)	184 (54.1%)	
Major	23 (11.0%)	54 (15.9%)	
Extreme	0	2 (0.6%)	

Symbols: \*:  $P < .05$ ; \*\*:  $P < .01$ ; \*\*\*:  $P < .001$ .

**TABLE 2. Comparison of Total Cost, Cost of Surgery, LOS, and Expected LOS Between NW and NICU**

	NW	NICU	Mann-Whitney P value
	Average; 95% CI		
Total cost	\$29 303.08; [27 613.00, 30 993.17]	\$32 496.41; [31 411.09, 33 581.73]	<.001***
Surgery cost	\$6994.29; [6812.62, 7175.97]	\$6833.52; [6698.91, 6968.13]	.327
Length of stay	3.046; [2.857, 3.235]	3.586; [3.439, 3.733]	<.001***
Expected length of stay	3.936; [3.829, 4.043]	3.889; [3.757, 4.022]	.184

Symbols: \*: P < .05; \*\*: P < .01; \*\*\*: P < .001.

**TABLE 3A. Univariate Linear Regression Analysis for LOS, Expected LOS, and Total Cost**

	Length of stay P value	Expected length of stay P value	Total cost P value
Admitting unit	<.001***	.628	.001**
Age at discharge	.201	.388	.102
MS DRG weight	<.001***	<.001***	<.001***
OR hours charged	<.001***	<.001***	0***
Sex	.318	.246	.891
Ethnicity	.921	<.001***	.070
Insurance	.080*	.218	.030*
Day of admission	<.001***	.047*	.470
Discharge disposition	0***	0***	0***
MS DRG	<.001***	<.001***	0***
APR DRG severity	<.001***	<.001***	<.001***

Symbols: \*: P < .05; \*\*: P < .01; \*\*\*: P < .001.

associated with longer LOS for the NW group, but not for the NICU group (Figure 2).

**Admitting to the NW Is Associated With Reduced Total Cost of Hospitalization**

We compared the average total cost of hospitalization between the NW and NICU groups. The NICU group averaged \$3193.33 more per admission (\$32 496.41 vs \$29 303.08; P < .001) (Table 2). There was no statistically significant difference in surgery cost between the 2 groups (\$6833.52 (NICU) vs \$6994.29 (NW); P = .327) (Table 2), suggesting that differences in costs incurred postoperatively.

To determine the specific cost components that contributed to the increased overall cost of the hospitalization, we compared 10 major components of cost between NW and NICU groups (Table 4A). The “room obs and overflow,” “room routine,” and “ICU room” categories correspond to the base cost of PACU, ward, and ICU equivalent levels of care, respectively. Notably, when examining the differences in component cost, the biggest difference was in the “ICU room” component, incurred by the

**TABLE 3B. Multivariate Linear Regression Analysis for LOS**


Included variables	Coefficient	Wald P value
Admitting unit (NICU)	0.235	.015*
MS DRG weight (larger)	−0.147	.028*
OR hours charged (more)	0.172	<.001***
Day of admission		<.001***
Monday	−	−
Tuesday	0.121	.362
Wednesday	0.196	.330
Thursday	0.681	<.001***
Friday	0.284	.071
Saturday	−0.373	.728
Discharge disposition		0***
Acute care	−	−
Deceased	1.434	.372
Home	−0.151	.779
Inpatient rehab	1.828	.001**
Skilled nursing facility	1.714	.006**
APR DRG severity		<.001***
Minor	−	−
Moderate	0.266	.023
Major	0.885	<.001***
Extreme	1.128	.296

Symbols: \*: P < .05; \*\*: P < .01; \*\*\*: P < .001.

NICU group. This difference in component cost is more than the increase the “room obs” and “room routine” components combined.

Multivariate analyses demonstrated that the admitting unit was not a significant independent predictor in the final model generated by stepwise regression. Variables independently associated with increased overall cost of hospitalization included OR hours charged, discharge disposition, MS-DRG, and APR-DRG severity (Table 4B). Interestingly, MS-DRG and APR-DRG severity were not significantly different between NW and NICU groups (Table 1).

In our post hoc analysis, however, the differences in cost between NW and NICU, when stratified by LOS, were the greatest when LOS was 1 d, and the effect diminishes with increasing LOS (Supplemental Digital Content 3, Table 3).



In multivariate analyses, NW admission was a significant independent predictor for total cost when LOS was less than 2, 3, 4, and 5 d (**Supplemental Digital Content 4-5, Tables 4-5**), independent of MS-DRG weight. This suggests that for LOS <6 d, NW admission becomes a strong determinant of total cost irrespective of intrinsic patient factors.

Further post hoc analyses demonstrated that boarding in PACU past midnight on postoperative day 0 was associated with increased overall cost for both the NW and NICU groups (Figure 3).

### NW and NICU Admissions Have Similar Clinical Outcomes

We evaluated whether admission to the NICU yielded better clinical outcome compared to the NW and found no statistically significant differences in rates of return to OR, return to Emergency Department (ED), or readmission within 30 d (Table 5).

Univariate logistical regression analyses revealed that sex was a predictor of return to OR within 30 d, while MS-DRG weight,

sex, and discharge disposition were predictors of readmission within 30 d (Table 6A). In multivariate analyses, male patients were found to be significantly more likely to return to the OR and be readmitted, while discharge disposition independently predicted whether a patient would be readmitted within 30 d (Table 6B). PACU boarding on postoperative day 0 was not associated with differential clinical outcomes for either cohort (**Supplemental Digital Content 6, Table 6**).

## DISCUSSION

The literature on the cost-effectiveness of neuroICU stay postoperatively for elective neurosurgical patients is scarce, in contrast to the existing literature on the benefits of NICU for brain and spinal cord injury patients.<sup>4-9</sup> Limited single institutional data from medical centers outside the United States have reported good outcomes with same day discharges for elective brain tumor craniotomy patients.<sup>10</sup> Despite the perceived clinical and economic benefits, however, outpatient craniotomies thus far have not gained widespread adoption.<sup>11</sup>

**TABLE 4A. Comparison of Cost Components Between NW and NICU**

	NW	NICU	Difference (NICU – NW)	Mann-Whitney P value
	Average; 95% CI			
Blood bank	\$166.95; [139.20, 194.70]	\$201.35; [169.69, 233.01]	\$34.40	.013*
ICU room	\$133.44; [51.89, 215.00]	\$2209.12; [2057.15, 2361.09]	\$2075.68	0***
Implant supplies	\$1806.61; [1477.36, 2135.85]	\$1896.93; [1705.52, 2088.35]	\$90.32	.459
Labs and pathology	\$1913.94; [1719.25, 2108.62]	\$2335.35; [2186.63, 2484.07]	\$421.41	<.001***
Pharmacy	\$2412.87; [1484.79, 3340.95]	\$2223.42; [1766.95, 2679.89]	–\$189.45	<.001***
Radiological imaging	\$299.28; [276.29, 322.27]	\$357.32; [330.44, 384.20]	\$58.04	<.001***
Radiological processing	\$201.10; [69.97, 332.22]	\$279.55; [163.47, 395.63]	\$78.45	.346
Room obs. and overflow	\$1020.05; [897.55, 1142.56]	\$538.22; [439.53, 636.91]	–\$481.83	<.001***
Room rehab	\$153.33; [120.47, 186.19]	\$227.15; [198.25, 256.05]	\$73.82	<.001***
Room routine	\$2706.91; [2465.81, 2948.02]	\$2310.14; [2090.27, 2530.02]	–\$396.77	<.001***

Symbols: \*:  $P < .05$ ; \*\*:  $P < .01$ ; \*\*\*:  $P < .001$ .

Room routine: cost of room and board in an NW equivalent room.

ICU room: cost of room and board in an ICU equivalent room.

**TABLE 4B. Multivariate Linear Regression Analysis for Total Costs**

Included variables	Coefficient	Wald P value
Admitting unit (NICU)	1047.30	.126
OR hours charged (more)	2402.60	0***
<b>Discharge disposition</b>		<.001***
Acute care	–	–
Deceased	18 309.90	.108
Home	1606.20	.675
Inpatient rehab	8931.20	.022*
Skilled nursing facility	9253.50	.035*
<b>MS DRG</b>		0***
23.00	–	–
25.00	–34 027.30	0***
26.00	–33 325.60	0***
27.00	–32 824.50	0***
41.00	–24 122.60	<.001***
42.00	–30 490.10	<.001***
54.00	–37 952.30	<.001***
55.00	–35 011.50	<.001***
<b>APR DRG severity</b>		<.001***
Minor	–	–
Moderate	2618.60	.004**
Major	6824.70	<.001***
Extreme	1880.50	.806

Symbols: \*:  $P < .05$ ; \*\*:  $P < .01$ ; \*\*\*:  $P < .001$ .


## Key Results

We present here a large retrospective analysis of a single academic institution's experience with elective supratentorial craniotomies for adult brain tumor patients without significant comorbidities with a LOS of less than 7 d. NICU and NW patients were well matched with respect to age, ethnicity, MS-DRG, APR-DRG, and expected LOS, with the exception of MS-

DRG weight, suggesting that most of the differences in outcome could be attributed to the relevant predicting variables we tested. Among those with LOS less than 5 d, even MS-DRG weight was well matched.

Our data suggest that even a single night stay in the NICU was associated with increased overall hospitalization cost and LOS, but with no difference in clinical outcome (return to ED, return to OR, and readmission within 30 d). The biggest contributor to this difference was the NICU's base cost. Our data imply that adult patients without significant comorbidities and with expected short LOS who undergo elective craniotomy for supratentorial lesions may be safely admitted to a NW unit without significant additional risks to the patient, while shortening their LOS and reducing hospitalization cost. Early recognition of this patient group and establishment of alternative pathways to ICU admission might decrease ICU utilization and the cost of hospitalization. NICU admission's independent effect on increased LOS could be explained by a variety of factors, including delayed mobilization, additional systemic barriers in multidisciplinary discharge planning, the lack of NW beds forcing a discharge directly from the NICU, and increased risk of ICU delirium.<sup>12</sup>

Our data also suggest that factors other than NICU admission can also contribute to longer LOS, given that the following variables were also independent predictors of LOS in our multivariate analysis: MS-DRG weight, OR hours charged, day of admission, discharge disposition, and APR-DRG severity. The longer surgery time (OR hours) and greater presence of comorbidities (APR-DRG severity) can explain a longer postoperative recovery time prior to discharge. Also, because discharging to a facility such as a skilled nursing facility or acute rehabilitation requires additional coordination and is dependent on the accepting facility's bed availability, it is not surprising that discharge disposition was a significant independent predictor of LOS.



Interestingly, NICU admission was not an independent predictor of higher cost in the multivariate model, unlike MS-DRG, APR-DRG severity, and discharge disposition. This suggests that the cost difference between the NW and the NICU groups could be better explained by differences in the presence of other comorbidities and where the patient was discharged than where they were admitted postoperatively. APR-DRG severity in particular as a potential confounder of NICU admission's effects on total cost implies that perhaps some neurosurgeons admitted patients who might benefit more from being in the ICU given their medical comorbidities, and the result of that was overall higher cost of hospitalization.

However, our subgroup analysis stratified by overall LOS found that the cost difference was greatest when LOS was short, and the effect diminishes as LOS lengthens. Because all patients in the NICU group only stayed in the NICU for 1 night, the diminishing effect with increasing LOS further highlights the need to identify patients who are otherwise healthy and expected to have shorter LOS, whose NW admission can potentially yield the greatest cost savings without affecting clinical outcome. This was supported by our finding that for the subgroup of patients with LOS less than 5 nights, NW admission independently predicted

lower total cost on multivariate analysis (**Supplemental Digital Content 4, Table 4**). Moreover, among those patients there were no significant differences in MS-DRG weight that would explain differences in cost due to the presence of higher level of comorbidities (**Supplemental Digital Content 5, Table 5**).

The subgroup analysis showing that extended PACU-stay past midnight increased LOS NW but not NICU admission can be due to the lack of a multidisciplinary team facilitating discharge in the PACU. The increased cost from extended PACU stay can be due to its higher staffing ratio, the anesthesia team, and higher resource utilization.

### Limitations

The limitations of our study include the retrospective nature of analysis, the lack of randomization to either the NW vs NICU group, the lack of multi-institutional data to reduce institutional bias, and the lack of more detailed data assessing the comorbidities of our patients which might affect their LOS. We also restricted our cohort of NICU patients to those who stayed in the NICU for 1 night. While this helped to ensure that most likely our NICU patient cohort had uneventful NICU recovery, it excluded those

**TABLE 5. Comparison of Clinical Outcomes Between NW and NICU**

	NW	NICU	
	Frequency		Pearson's chi-squared P-value
<b>Return to OR</b>			.867
Yes	8 (3.8%)	14 (4.1%)	
No	201 (96.2%)	326 (95.9%)	
<b>Return to ED</b>			.757
Yes	21 (10.0%)	37 (10.9%)	
No	188 (90.0%)	303 (89.1%)	
<b>Readmission</b>			.398
Yes	22 (10.5%)	44 (12.9%)	
No	187 (89.5%)	296 (87.1%)	

Symbols: \*:  $P < .05$ ; \*\*:  $P < .01$ ; \*\*\*:  $P < .001$ .

**TABLE 6A. Univariate Logistic Regression Analysis for Return to OR, Return to ED, and Readmission Within 30 Days**

	Return to OR P value	Return to ED P value	Readmission P value
Admitting unit	.866	.757	.399
Age at discharge	.592	.534	.498
MS DRG weight	.908	.514	.009**
OR hours charged	.970	.879	.409
Sex	.041*	.651	.006**
Ethnicity	.998	.380	.998
Insurance	.067	.560	.800
Day of admission	.870	.950	.380
Discharge disposition	.390	.850	<.001***
MS DRG	.580	.920	.062
APR DRG severity	.950	.730	.880

Symbols: \*:  $P < .05$ ; \*\*:  $P < .01$ ; \*\*\*:  $P < .001$ .

**TABLE 6B. Multivariate Logistic Regression Analysis for Return to OR and Readmission Within 30 Days**

Included variables	Odds ratio	Wald P value
<b>Return to OR</b>		
Sex (male)	2.697	.041*
<b>Readmission</b>		
Sex (male)	2.190	.008**
<b>Discharge disposition</b>		
Acute care	–	–
Deceased	0	.988
Home	0.299	.305
Inpatient rehab	2.607	.418
Skilled nursing facility	1.424	.793

Symbols: \*:  $P < .05$ ; \*\*:  $P < .01$ ; \*\*\*:  $P < .001$ .

who stayed longer in the NICU but also had uneventful recoveries. Consequently, we may have artificially reduced the degree of differences that can be seen between NW and NICU group analyses both in terms of LOS and from a cost perspective. It would not be surprising to see a trend toward continued increase in LOS and hospitalization cost by including those who stayed in the NICU more than 1 night, on a per nightly basis.

### Interpretation and Generalizability

Our analysis demonstrated that even a 1-night stay in the NICU was sufficient to see an increase in costs. A potential limitation of our study is its generalizability given our practice setting of an academic center. In our university hospital, the NW is staffed with nurses who have undergone rigorous training under a preceptor and a 6-mo probationary period prior to independent patient care. Additionally, the nurse to patient ratio is either 1:3 or 1:4 allowing for neurological assessments every 2 h. This in conjunction with telemetry monitoring allows for the NW to serve as a neurological surgery stepdown unit in a way that may not be replicable at other centers. However, given the appropriate institutional setting, our study supports the creation of postoperative pathways that allow for select patients without significant medical comorbidities for admission to a neurosurgical ward.

### CONCLUSION

Admitting adult elective craniotomy patients without significant medical comorbidities who are expected to have a short LOS to the NW is independently associated with reduced LOS compared to admitting to the NICU for 1 night prior to the NW. The severity of comorbidities and discharge disposition are independent predictors of total cost. In patients with LOS < 5 d, direct postoperative NW admission may even be an independent predictor of reduced hospitalization costs, without significant differences in clinical outcomes.

### Funding

This study did not receive any funding or financial support.

### Disclosures

The authors have no personal, financial, or institutional interest in any of the drugs, materials, or devices described in this article.

### REFERENCES

- Hunter A, Johnson L, Coustasse A. Reduction of intensive care unit length of stay: the case of early mobilization. *Health Care Manag.* 2014;33(2):128-135.
- Kramer AA, Zimmerman JE. The relationship between hospital and intensive care unit length of stay. *Crit Care Med.* 2011;39(5):1015-1022.
- Raj R, Bendel S, Reinikainen M, et al. Costs, outcome and cost-effectiveness of neurocritical care: a multi-center observational study. *Crit Care.* 2018;22(1):225.
- Diringner MN, Edwards DF. Admission to a neurologic/neurosurgical intensive care unit is associated with reduced mortality rate after intracerebral hemorrhage. *Crit Care Med.* 2001;29(3):635-640.
- Mirski MA, Chang CW, Cowan R. Impact of a neuroscience intensive care unit on neurosurgical patient outcomes and cost of care: evidence-based support



- for an intensivist-directed specialty ICU model of care. *J Neurosurg Anesthesiol.* 2001;13(2):83-92.
6. Varelas PN, Conti MM, Spanaki MV, et al. The impact of a neurointensivist-led team on a semiclosed neurosciences intensive care unit. *Crit Care Med.* 2004;32(11):2191-2198.
  7. Suarez JL, Zaidat OO, Suri MF, et al. Length of stay and mortality in neurocritically ill patients: impact of a specialized neurocritical care team. *Crit Care Med.* 2004;32(11):2311-2317.
  8. Varelas PN, Eastwood D, Yun HJ, et al. Impact of a neurointensivist on outcomes in patients with head trauma treated in a neurosciences intensive care unit. *J Neurosurg.* 2006;104(5):713-719.
  9. Bershad EM, Feen ES, Hernandez OH, Suri MF, Suarez JL. Impact of a specialized neurointensive care team on outcomes of critically ill acute ischemic stroke patients. *Neurocrit Care.* 2008;9(3):287-292.
  10. Purzner T, Purzner J, Massicotte EM, Bernstein M. Outpatient brain tumor surgery and spinal decompression: a prospective study of 1003 patients. *Neurosurgery.* 2011;69(1):119-127; discussion 126-117.
  11. Marigil M, Bernstein M. Outpatient neurosurgery in neuro-oncology. *Neurosurg Focus.* 2018;44(6):E19.
  12. Herling SF, Greve IE, Vasilevskis EE, et al. Interventions for preventing intensive care unit delirium in adults. *Cochrane Database Syst Rev.* 2018;11(11):CD009783.

---

**Supplemental digital content** is available for this article at [www.neurosurgery-online.com](http://www.neurosurgery-online.com).

**Supplemental Digital Content 1. Table 1:** List of ICD codes.

**Supplemental Digital Content 2. Table 2:** List of variables.

**Supplemental Digital Content 3. Table 3:** Mean total cost between NW and NICU stratified by total LOS.

**Supplemental Digital Content 4. Table 4:** NW admission was an independent predictor of total cost in shorter LOS subgroups.

**Supplemental Digital Content 5. Table 5:** Mean MS-DRG weight between NW and NICU in shorter LOS subgroups.

**Supplemental Digital Content 6. Table 6:** Comparison of clinical outcomes stratified by extended PACU stay. "PACU to NW" and "PACU to NICU" groups stayed in the PACU past midnight on postoperative day 0. Patients who were transferred from the OR directly to the NICU were in the "directly to NICU" group. Patients who were admitted to the NW from PACU within a few hours of surgery and did not stay past midnight were in the "directly to NW" group. Symbols: \*:  $P < .05$ ; \*\*:  $P < .01$ ; \*\*\*:  $P < .001$ .

---